## 120. COASTAL PRAIRIE

PRADERA COSTANERA

JOHN P. KELLY

Audubon Cypress Grove Preserve P.O. Box 808 Marshall, California 94940

Location: California; Marin Co.; Marshall; Cypress Grove Preserve; 38°10′N, 122°54′W; Tomales Quadrangle, USGS. Continuity: Established 1988; 6 yr. Size: 31.0 ha. Description of Plot: See J. Field Ornithol. 60 (Suppl.):56–57 (1989). Detailed quantitative habitat data from 1993 are available from the author. Weather: Mean start temp., 13.0°C (range 8–17°C). Generally mild with normal temperatures. Rainfall during the 1992–93 rainy season was slightly (20%) above the 38-yr average of 71.9 cm, following six consecutive years of below-average rainfall. Significant periods of rainfall extended into April and May. Source: Audubon Can-

yon Ranch. Coverage: 7.7 h; 8 visits (2 sunrise); 13, 27 April; 12, 19, 21, 25, 27, 28 May. Census: Song Sparrow, 16.0 (21); Black-shouldered Kite, 2.0; Northern Harrier, 2.0; Mourning Dove, 2.0; American Goldfinch, 2.0; Red-winged Blackbird, 1.5; American Crow, 1.0; California Towhee, 1.0; Western Meadowlark, 1.0; Allen's Hummingbird, 0.5; Northern Oriole, +. Total: 11 species; 29.0 territories (37/40 ha). Visitors: Mallard, Osprey, Anna's Hummingbird, Western Kingbird, Cliff Swallow, Barn Swallow, Scrub Jay, Bushtit, Marsh Wren, Tricolored Blackbird, Brewer's Blackbird, Brown-headed Cowbird, House Finch, Pine Siskin. Remarks: At least one of the two Black-shouldered Kite nests successfully fledged young. An increase of one Northern Harrier territory occurred this year as vole (Microtus californicus) densities increased over levels observed since 1989–90. Acknowledgments: This work is supported by Audubon Canyon Ranch, Stinson Beach, CA. I thank Susan Kelly and Sarah Tappen for help in summarizing the results.